

Pain after spine surgery

Why pain can persist, return, or change after an operation.

What does this mean?

Pain may continue after decompression, disc surgery, or fusion, or return after an initial improvement. Terms such as post-laminectomy syndrome describe the situation, not one specific pain source.




Illustration of lumbar fusion: front and side views.

Persistent pain deserves a fresh evaluation.

It does not automatically mean the operation failed or that another surgery is needed.

Nerve-related pain

Burning, electric, tingling, or shooting pain may travel into an arm or leg. A nerve can remain sensitive after pressure is relieved, or may be affected by new narrowing or scar tissue.

Joint, disc, or muscle pain

Facet joints, sacroiliac joints, nearby discs, and supporting muscles may contribute. Changes in movement after fusion can place more demand on nearby structures.

Structural causes

Your team may assess for recurrent narrowing, a new disc problem, incomplete fusion, instability, or a hardware problem. Findings need to match the symptoms before treatment is chosen.

More than one source

Back pain and leg pain may have different causes. Treating nerve pain alone may leave joint or muscle pain untreated. The symptom pattern guides the next step.

Pain can affect the whole recovery

Poor sleep, reduced activity, and fear of movement can add to persistent pain. Addressing these factors is part of care and does not mean your pain is imagined.

Finding a path forward

Match the treatment to the cause, then measure what improves.

Start with a careful review

Bring your operation report, previous scans, medication list, and details of treatments you have tried. Explain what improved after surgery, what did not, and when symptoms changed.

Choose tests for a reason

The examination checks movement, strength, sensation, and walking. Depending on the question, your team may use MRI, CT, X-rays, or nerve testing. A diagnostic injection may help identify a joint-related pain source.

Rebuild activity gradually

A tailored rehabilitation plan can improve strength, endurance, and confidence. Medicines may help nerve pain or make activity more manageable. Sleep support and pain-coping skills can also help.

Targeted procedures

Depending on the diagnosis, options may include an epidural injection, a joint injection, or radiofrequency treatment after an appropriate diagnostic block. Procedures do not address every cause of pain.

Spinal cord stimulation

For selected patients with persistent pain despite other care, stimulation may reduce pain signaling. A temporary trial usually helps assess benefit before a permanent implant. It does not repair the spine. Risks include infection, bleeding, lead movement, device problems, and incomplete relief.

When is more surgery useful?

Another operation may be appropriate when there is a clear, treatable structural problem that fits your symptoms. Persistent pain by itself is not enough to decide. Review the likely benefit and risks with your surgeon.

What would meaningful progress look like?

Choose two or three goals, such as walking farther, sleeping better, sitting comfortably, or returning to work. Ask: Which pain source are we treating? How will we judge benefit? What is the next step if it does not help?



Reza S. Roghani, MD, PhD

Assistant Professor
Interventional Spine and Pain

Department of Physical Medicine and Rehabilitation
School of Medicine

443-997-5476 Appointments
410-614-4033 Fax
410-502-2199 Procedures

Appointments

443-997-5476

Johns Hopkins Medicine

General education; individual evaluation is required.